# Ca<sub>v</sub>1.3 Is Preferentially Coupled to Glucose-Stimulated Insulin Secretion in the Pancreatic $\beta$ -Cell Line INS-1

GUOHONG LIU, NEJMI DILMAC, NATHAN HILLIARD, and GREGORY H. HOCKERMAN

Department of Medicinal Chemistry and Molecular Pharmacology (G.L., N.D., N.H., G.H.H.) and Neuroscience Graduate Program (G.L.), Purdue University, West Lafayette, Indiana

Received October 30, 2002; accepted December 30, 2002

## ABSTRACT

L-Type Ca<sup>2+</sup> channel blockers inhibit glucose and KCI-stimulated insulin secretion by pancreatic  $\beta$  cells. However, the role of the two distinct L-type channels expressed by  $\beta$  cells, Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3, in this process is not clear. Therefore, we stably transfected INS-1 cells with two mutant channel constructs, Ca<sub>v</sub>1.2DHPi or Ca<sub>v</sub>1.3 DHPi. Whole-cell patch-clamp recordings demonstrated that both mutant channels are insensitive to dihydropyridines (DHPs), but are blocked by diltiazem. INS-1 cells expressing Ca<sub>v</sub>1.3/DHPi maintained glucose- and KCI-stimulated insulin secretion in the presence of DHPs, whereas cells expressing Ca<sub>v</sub>1.2/DHPi demonstrated DHP resistance to only KCI-induced secretion. INS-1 cells were also stably trans-

fected with cDNAs encoding the intracellular loop between domains II and III of either Ca<sub>v</sub>1.2 or Ca<sub>v</sub>1.3 (Ca<sub>v</sub>1.2/II-III or Ca<sub>v</sub>1.3/II-III). Glucose- and KCI-stimulated insulin secretion in Ca<sub>v</sub>1.2/II-III cells were not different from untransfected INS-1 cells. However, glucose-stimulated insulin secretion was completely inhibited and KCI-stimulated secretion was substantially resistant to inhibition by DHPs, but sensitive to  $\omega$ -agatoxin IVA in Ca<sub>v</sub>1.3/II-III cells. Moreover, the L-type channel agonist FPL 64176 markedly enhanced KCI-stimulated secretion by Ca<sub>v</sub>1.3/II-III cells. Together, our results suggest that Ca<sup>2+</sup> influx via Ca<sub>v</sub>1.3 is preferentially coupled to glucose-stimulated insulin secretion in INS-1 cells.

Calcium influx through voltage-dependent Ca<sup>2+</sup> channels (VDCCs) plays a crucial role in insulin secretion from pancreatic  $\beta$  cells (Wollheim and Sharp, 1981). Glucose metabolism in  $\beta$  cells causes an increase in ATP/ADP ratio, which closes ATP-dependent potassium channels (Rajan et al., 1990). The resulting membrane depolarization opens VD-CCs, and calcium influx triggers insulin secretion. Several distinct VDCCs have been detected in  $\beta$  cells and insulinsecreting cell lines, including high-voltage-activated subtypes (Seino et al., 1992; Ligon et al., 1998) and the lowvoltage-activated channels (Zhuang et al., 2000). L-Type VDCCs play a major role in the function of pancreatic  $\beta$  cells because L-type-specific blockers significantly inhibit glucose or depolarization-induced insulin secretion (Devis et al., 1975). Two distinct L-type channels, Ca, 1.2 and Ca, 1.3, are expressed in human pancreatic islets (Seino et al., 1992) and in insulin-secreting cells lines (Horvath et al., 1998). However, Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3 are both blocked by the same smallmolecule drugs (Hockerman et al., 1997; Bell et al., 2001). Thus, the relative contribution of calcium flux through  $\rm Ca_v 1.2$  and/or  $\rm Ca_v 1.3$  to  $\rm Ca^{2+}$ -dependent insulin secretion is not clear.

Previously, we reported a double mutation that renders  $Ca_v1.2$  insensitive to dihydropyridines (DHPs), such as nifedipine and PN200-110, but normally sensitive to block by diltiazem (Hockerman et al., 2000). We used this mutant channel ( $Ca_v1.2$ /DHPi), and the corresponding  $Ca_v1.3$  mutant ( $Ca_v1.3$ /DHPi), in a novel strategy to study the roles of  $Ca_v1.2$  (Snutch et al., 1991) and  $Ca_v1.3$  (Williams et al., 1992) in insulin secretion. When these mutant channels were introduced into the rat pancreatic  $\beta$ -cell line INS-1 (Asfari et al., 1992), endogenous L-type channels were "turned off" with a DHP such as PN200-110, functionally isolating the drug-insensitive  $Ca_v1.2$  or  $Ca_v1.3$  mutant. We found that expression of  $Ca_v1.3$ /DHPi but not  $Ca_v1.2$ /DHPi allowed glucose-stimulated insulin secretion that was insensitive to DHPs, but blocked by diltiazem, in INS-1 cells.

Insulin-containing secretory granules and L-type VDCCs are colocalized in  $\beta$ -cells (Bokvist et al., 1995; Qian and Kennedy, 2001). The functional coupling between L-type channels and exocytotic granules may resemble that of non-

This work was supported by Grant 1119990378 from the American Diabetes Association (to G.H.H.).

Article, publication date, and citation information can be found at http://jpet.aspetjournals.org.

DOI: 10.1124/jpet.102.046334.

**ABBREVIATIONS:** VDCC, voltage-dependent calcium channel; DHP, dihydropyridine; DHPi, dihyrdropyridine-insensitive; SNARE, soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor; GFP, green fluorescent protein; RT-PCR, reverse transcription-polymerase chain reaction; EGFP, enhanced green fluorescent protein; PCR, polymerase chain reaction; bp, base pair(s); KRBH, Krebs-Ringer-bicarbonate HEPES buffer; *I*<sub>Ba</sub>, barium current.

L-type VDCCs to neurotransmitter release in neurons. An interaction between the intracellular loop linking domains II and III of Ca<sub>v</sub>2.2 or Ca<sub>v</sub>2.1 and SNARE proteins mediates efficient coupling of Ca<sup>2+</sup> influx to neurotransmitter vesicle fusion (Sheng et al., 1998). Pancreatic  $\beta$ -cells express isoforms of the corresponding SNARE proteins, which regulate both insulin secretion and calcium channel activity (Jacobsson et al., 1994; Sadoul et al., 1995; Nagamatsu et al., 1996). Functional interactions between syntaxin and both Ca<sub>v</sub>1.3 (Yang et al., 1999) and Ca<sub>v</sub>1.2 (specifically the II-III loop) (Wiser et al., 1999) are reported to play a role in depolarization-induced insulin secretion. To examine the role of the II-III loop of both Ca, 1.2 and Ca, 1.3 in coupling Ca<sup>2+</sup> influx to insulin secretion, we stably transfected INS-1 cells with plasmids encoding the II-III loop of either Ca<sub>v</sub>1.2 or Ca<sub>v</sub>1.3, fused to GFP. Experiments with the resulting cell lines (Cav1.2/II-III cells and Cav1.3/II-III cells) showed that glucose-stimulated insulin secretion was completely abolished in Ca, 1.3/II-III cells, but was not different from untransfected cells in the Ca<sub>v</sub>1.2/II-III cell line.

# **Materials and Methods**

**Cell Culture.** INS-1 cells were cultured as reported previously (Asfari et al., 1992).

**Stable Transfection.** INS-1 cells were transfected using Gene-PorterII (Gene Therapy Systems, San Diego, CA). After 3 days, 100  $\mu$ g/ml G418 (Promega, Madison, WI) was added to the medium. Colonies were isolated, and subsequently screened by RT-PCR and Western blot.

**Plasmids.** Construction of the Ca<sub>v</sub>1.2/DHPi mutant was described previously (Hockerman et al., 2000). The two amino acid mutations in domain IIIS5 of Ca<sub>v</sub>1.3 (Williams et al., 1992) (Thr 1029 to Tyr; Gln 1033 to Met) were mutated using the QuikChange method (Stratagene, La Jolla, CA) to generate Ca<sub>v</sub>1.3/DHPi. Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi were subcloned into EGFP vector (BD Biosciences Clontech, Palo Alto, CA). The intracellular II-III loops of Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3 were amplified by PCR using Pfu DNA polymerase, followed by ligation into the EGFP vector. All constructs were confirmed by cDNA sequencing.

RT-PCR. Total RNA was extracted from INS-1 cells using TRIzol (Invitrogen, Carlsbad, CA), and 2  $\mu$ g were incubated with random primers at 70°C for 5 min, and then put on ice. RNase inhibitor  $(1 \mu l)$ , 100 µM dNTPs, 0.01 M dithiothreitol, and 200 U of M-MLV reverse transcriptase (Promega, Madison, WI) were added to the mixture (20  $\mu$ l final volume) and incubated at 42°C for 30 min, followed by incubation at 85°C for 5 min. Two primer pairs were used for PCR with Taq polymerase (Promega) as follows. Mutant primer set (overlaps mutations in IIIS5): Ca, 1.2 forward (5'-cta cac tct gct gat gtt c-3') and reverse (5'-ggg gat cca cgt acc aca ctt tgt act-3') and Ca<sub>v</sub>1.3 forward (5'-cat gac cct cct gat gtt c-3') and reverse (5'-cgg gat ccc gcg aag agt tca cca cgt ac-3'). PCR products are 538 bp for Ca<sub>v</sub>1.2/DHPi and 540 bp for Ca<sub>v</sub>1.3/DHPi. GFP primers set: Ca<sub>v</sub>1.2 (5'-agc tgt gta tat gcc ctg g-3'), GFPr (5'-gaa gaa gtc gtg ctg ctt c-3'), Ca<sub>v</sub>1.3 (5'-gtc ctg gct aca gcg acg-3'), and GFPr were used to amplify the channel/ EGFP junction. PCR products are 344 bp for Ca, 1.2/DHPi and 351 bp for Ca<sub>v</sub>1.3/DHPi. PCR products were visualized by ethidium bromide staining after 1% agarose gel electrophoresis in TAE buffer (40 mM Tris-acetate, 2 mM EDTA, pH 8.5).

Western Blot. Crude membranes from indicated cells were isolated as described previously (Peterson et al., 1997). For whole-cell lysates, indicated cells were incubated in SDS lysis buffer (0.5% SDS, 0.05 M Tris-Cl, 1 mM dithiothreitol, pH 8.0) for 10 min. Lysates were boiled for 5 min and clarified by centrifugation at 26,000g, 4°C for 90 min, and supernatants were collected for Western blot. The proteins were separated by SDS-polyacrylamide gel electrophoresis (5% gels for crude membranes and 12% gels for cell lysates) followed by transfer to nitrocellulose membrane. The membranes were blocked with 5% nonfat milk in Tris-buffered saline at 4°C overnight, washed with 0.1% Tween 20 in Tris-buffered saline, and incubated with polyclonal rabbit anti-GFP antibodies (BD Biosciences Clontech) for 2 to 3 h. The blots were detected by incubation with horseradish peroxidase-conjugated anti-rabbit antibodies and visualized by enhanced chemiluminescence with Hyperfilm (Amersham Pharmacia AB, Uppsala, Sweden). Protein concentrations were determined using the Bradford assay (Bio-Rad, Hercules, CA).

Insulin Secretion and Cellular Insulin Content. Cells in 12well dishes were rinsed twice with modified Krebs-Ringer-bicarbonate HEPES buffer (KRBH buffer: 115 mM NaCl, 24 mM NaHCO<sub>3</sub>, 5 mM KCl, 1 mM MgCl<sub>2</sub>, 2.5 mM CaCl<sub>2</sub>, 25 mM HEPES, 0.5% BSA, pH 7.4), and incubated in 1 ml KRBH buffer for 30 min at 37°C, 5% CO<sub>2</sub>. After aspiration of the buffer, the cells were stimulated in KRBH buffer supplemented with 50 mM KCl or 11.2 mM glucose (±indicated drugs) for 30 min at 37°C. The buffer containing secreted insulin was collected and centrifuged at 700g for 3 min to remove any detached cells. The cellular insulin was extracted by incubation in 1 ml of acid-ethanol (ethanol/H<sub>2</sub>O/HCl, 14:57:3) overnight at 4°C and centrifuged at 700g for 3 min. Supernatants were diluted 10-fold with glycine-NaOH buffer (0.2 M glycine, 0.5% bovine serum albumin, pH 8.8) before insulin assay. Assays were performed using the High Range Rat Insulin ELISA kit (ALPCO, Windham, NH) according to the manufacturer's instructions.

**Electrophysiology.** Cells were cultured on plastic coverslips for 2 days (Nalge Nunc, Naperville, IL). Whole-cell barium currents were recorded at room temperature using an Axopatch 200B amplifier (Axon Instruments, Inc., Foster City, CA) and filtered at 1 kHz (six-pole Bessel filter, -3 dB). Electrodes were pulled from borosilicate glass (VWR, West Chester, PA) and fire polished to resistances of 2 to 6 MΩ. Voltage pulses were applied and data were acquired using pClamp8 software (Axon Instruments, Inc.). Nifedipine and diltiazem (Sigma-Aldrich, St. Louis, MO) were applied to the recording chamber in bath saline at 0.5 ml/min. The bath saline contained 150 mM Tris, 10 mM BaCl<sub>2</sub>, and 4 mM MgCl<sub>2</sub>. The intracellular solution contained 130 mM *N*-methyl-D-glucamine, 10 mM EGTA, 60 mM HEPES, 2 mM MgATP, and 1 mM MgCl<sub>2</sub>. The pH of both solutions was adjusted to 7.3 with methanesulfonic acid.

**Data Analysis.** Data were analyzed using ClampFit 8.1 (Axon Instruments, Inc.) and SigmaPlot 2001 (SPSS Science, Chicago, IL). Data are shown as mean values  $\pm$  standard error. Statistical significance was determined using Student's *t* test.

## **Results**

Molecular Characterization of Ca<sub>v</sub>1.2/DHPi and Cav1.3/DHPi Stable Cell Lines. The Cav1.2/DHPi and Ca<sub>v</sub>1.3/DHPi mutant constructs incorporate Tyr for Thr and Met for Gln substitutions in IIIS5 that render them insensitive to DHPs, but do not affect sensitivity to diltiazem block (Hockerman et al., 2000) (Fig. 1A). INS-1 cells were transfected with either Ca<sub>v</sub>1.2/DHPi or Ca<sub>v</sub>1.3/DHPi fused to GFP. After selection in G418, colonies were screened by RT-PCR. Because both Ca, 1.2 and Ca, 1.3 are endogenously expressed in INS-1 cells, we used oligonucleotide primers complementary to the mutations in transmembrane domain IIIS5, or primers complementary to GFP cDNA in the PCR reactions. As shown in Fig. 1B, RT-PCR reactions using RNA from two different clones from each transfection, and both primer pairs, amplified DNA fragments of the expected size. Control reactions using the same primers with RNA from untransfected INS-1 cells did not amplify the corresponding DNA fragments. We further characterized one clone from each transfection by Western blot. Because antibodies to Ca, 1.2 or



**Fig. 1.** Stable expression of Ca<sub>v</sub>1.2/DHPi or Ca<sub>v</sub>1.3/DHPi in INS-1 cells. A, amino acids mutated in Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi. Both constructs included GFP fused to the C-terminal tail. B, RT-PCR of stable INS-1 cells expressing Ca<sub>v</sub>1.2/DHPi or Ca<sub>v</sub>1.3/DHPi. Primers complementary to the IIIS5 mutations or GFP were used as described under *Materials and Methods*. RNA from untransfected INS-1 cells served as control. C, Western blot analysis. Crude membrane fractions from untransfected INS-1 cells and selected cell lines subjected to SDS-polyacryl-amide gel electrophoresis and blotted with anti-GFP antibodies. A protein of ~240 kDa was detected in both stable cell lines, but not in untransfected INS-1 cells.

 $Ca_v 1.3$  would detect endogenous channels, we used an anti-GFP antibody to detect the mutant channels. Figure 1C shows that anti-GFP antibodies detect a protein migrating with a molecular mass of approximately 240 kDa in crude membrane fractions from  $Ca_v 1.2$ /DHPi cells, and a slightly larger protein in crude membranes prepared from  $Ca_v 1.3$ / DHPi cells. Neither protein was detected by anti-GFP antibodies in crude membrane fractions from untransfected INS-1 cells.

**Electrophysiological Characterization of Ca**<sub>v</sub>**1.2**/ **DHPi and Ca**<sub>v</sub>**1.3**/**DHPi Cell Lines.** Both Ca<sub>v</sub>**1.2**/DHPi and Ca, 1.3/DHPi cell lines were characterized using whole-cell patch-clamp recordings of  $Ba^{2+}$  currents ( $I_{Ba}$ ). Stable expression of the  $Ca_v 1.2/DHPi$  or  $Ca_v 1.3/DHPi$  channels did not significantly increase the  $I_{\rm Ba}$  density compared with untransfected INS-1 cells (Fig. 2B). Both 1  $\mu$ M PN200-110 and 10  $\mu$ M nifedipine block approximately 13% of  $I_{\rm Ba}$  in untransfected INS-1 cells, and coapplication of 50  $\mu$ M diltiazem with 10  $\mu$ M nifedipine does not significantly reduce current amplitude (Fig. 2, A and C). The fraction of current blocked by application of 10 µM nifedipine to Ca, 1.2/DHPi cells or Ca, 1.3/DHPi cells under voltage clamp (Fig. 2, A and C) was not different from untransfected INS-1 cells. However, coapplication of 50  $\mu$ M diltiazem along with 10  $\mu$ M nifedipine to Ca<sub>v</sub>1.2/DHPi cells or Ca<sub>v</sub>1.3/DHPi cells further reduced  $I_{\rm Ba}$ . Thus, the pharmacological profile of  $I_{\rm Ba}$  in Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/ DHPi cells confirms the functional expression of the mutant channels. Furthermore, the  $I_{Ba}$  density (pA/pF) is not significantly different in the Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cells lines compared with untransfected INS-1 cells (Fig. 2B). Thus, stable transection of INS-1 cells with the mutant  $\alpha_1$ subunits alone, does not lead to a significant increase in whole-cell barium current.

Insulin Secretion in Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi Cells. We tested the ability of the Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/ DHPi cells to secretion insulin in response to either KCl or glucose stimulation, in the presence or absence of a DHP channel blocker. Glucose- and KCl-induced insulin secretion in untransfected INS-1 cells is completely blocked by the DHP PN200-110 (0.1 µM) (Fig. 3A). Glucose-stimulated insulin secretion in Ca<sub>v</sub>1.2/DHPi cells is completely blocked by 1  $\mu$ M PN200-110 or 1  $\mu$ M PN200-110 + 500  $\mu$ M diltiazem (Fig. 3B). However, glucose-stimulated insulin secretion in  $Ca_v 1.3$ /DHPi cells is substantially resistant to 1  $\mu$ M PN200-110, but inhibited by the addition of 500  $\mu$ M diltiazem. In contrast, KCl-induced insulin secretion is resistant to PN200-110 in both Ca<sub>v</sub>1.3/DHPi and Ca<sub>v</sub>1.2/DHPi cells (Fig. 3C). Figure 3D summarizes the results of these experiments by comparing the percentage of glucose- and KCl-stimulated insulin secretion resistant to 1  $\mu$ M PN200-110 in both Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cells. In Ca<sub>v</sub>1.3/DHPi cells,  $\sim$ 42% of glucose-stimulated insulin secretion was resistant to PN200-100, whereas glucose did not stimulate insulin secretion from Ca. 1.2/DHPi cells in the presence of PN200-110. Upon KCl stimulation,  $\sim 44\%$  of insulin secretion was resistant to PN200-110 in Ca<sub>v</sub>1.3/DHPi cells, whereas a significantly lower percentage of insulin secretion ( $\sim 29\%$ ) was resistant to PN200-110 in Ca, 1.2/DHPi cells. Thus, although the percentage of DHP-resistant insulin secretion was the same regardless of the stimulus in Ca, 1.3/DHPi cells, DHPresistant insulin secretion in Ca<sub>v</sub>1.2/DHPi cells was dependent upon KCl stimulation. The portion of DHP-sensitive secretion in both cell lines suggests that some endogenous L-type channels remain functionally coupled to secretion.

Characterization of Ca<sub>v</sub>1.2/II-III and Ca<sub>v</sub>1.3/II-III Cells. The intracellular II-III loops of several Ca<sub>v</sub> channels are known to bind to other signaling proteins, thus efficiently coupling Ca<sup>2+</sup> influx to a cellular response (Sheng et al., 1994; Rettig et al., 1996; Grabner et al., 1999). Therefore, we hypothesized that overexpression of the Ca<sub>v</sub>1.3 II-III loop should inhibit glucose-stimulated insulin secretion if an interaction between this domain and another protein is critical for this specific signaling pathway. To test this hypothesis,



Fig. 2. Electrophysiological characterization of Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cells.  $I_{Ba}$  was measured using whole-cell voltage-clamp. Currents were elicited by 100-ms depolarizations from -60 to +10 mV. A, representative traces from untransfected INS-1 cells, Ca<sub>v</sub>1.2/DHPi cells, and Ca<sub>v</sub>1.3/DHPi cells. Nifedipine (Nif; 10  $\mu$ M) blocked DHP-sensitive current. Nifedipine (10  $\mu$ M) plus diltiazem (dilt; 50  $\mu$ M) blocked Ca<sub>v</sub>1.2/DHPi or Ca<sub>v</sub>1.3/DHPi cells. B, barium current density in INS-1,Ca<sub>v</sub>1.2/DHPi, and Ca<sub>v</sub>1.3/DHPi cells. Total whole-cell current (pA/pF) measured (mean ± S.E.; n = 4-6) is shown. There is no statistically significant difference among the three cell lines tested. C, relative fractional blockade of  $I_{Ba}$  by drugs. In untransfected INS-1 cells, 13.5 ± 3% (n = 4) of total barium current was blocked by 10  $\mu$ M nifedipine, whereas 12 ± 2.5% (n = 6) was blocked by 1  $\mu$ M PN200-110. Coapplication of 50  $\mu$ M diltiazem did not significantly reduce DHP-resistant current. In Ca<sub>v</sub>1.2/DHPi cells, ca<sub>v</sub>1.3/DHPi cells, p < 0.05].

we created stable INS-1 cell lines that express the intracellular II-III loop of either  $Ca_v 1.2$  or  $Ca_v 1.3$  fused via the C terminus to GFP ( $Ca_v 1.2$ /II-III and  $Ca_v 1.3$ /II-III) (Fig. 4A). Western blot analysis of  $Ca_v 1.2$ /II-III and  $Ca_v 1.3$ /II-III cells using anti-GFP antibodies detected proteins of ~43 and ~45 kDa, respectively, consistent with the expected molecular mass of each fusion protein (Fig. 4B). Neither protein was detected in untransfected INS-1 cells.

Insulin Secretion in Ca, 1.3/II-III and Ca, 1.2II-III Cells. Fig. 4C shows that glucose-stimulated insulin secretion was maintained and completely inhibited by PN200-110 in Ca. 1.2/II-III cells. However, in Ca. 1.3/II-III cells, glucose did not stimulate insulin secretion. Upon KCl stimulation (Fig. 4D), Cav1.2/II-III cells again demonstrated normal insulin secretion that was completely blocked by PN200-110. However, KCl-stimulated insulin secretion by Ca, 1.3/II-III cells was sharply decreased compared with untransfected INS-1 or Ca<sub>v</sub>1.2/II-III cells, and it was substantially resistant to block by PN200-110. Analysis of  $I_{\rm Ba}$  in Ca<sub>v</sub>1.3/II-III cells detected a normal level of L-type channel activity (data not shown). Thus, we asked whether enhancing the  $Ca^{2+}$  influx via the endogenous L-type channels could increase KCl-stimulated secretion. Figure 5A shows the effect of the L-type channel agonist FPL 64176 on KCl-stimulated insulin secretion in Ca<sub>v</sub>1.3/II-III cells. As before, KCl evoked a modest level of insulin secretion, which was not sensitive to PN200-110. When FPL 64176 (10  $\mu$ M) was applied to Ca<sub>.</sub>1.3/II-III cells, KCl-stimulated secretion was sharply increased (>440%; Fig. 5A, inset). This increase was completely blocked by 1  $\mu$ M PN200-110. In contrast, when FPL 64167 is applied to untransfected INS-1 cells, insulin secretion is increased by only 174% (Fig. 5A, inset). These data suggest that overexpression of the Ca<sub>v</sub>1.3 II-III loop may spatially separate the endogenous Ca<sub>v</sub>1.3 channels from the cell's secretory machinery. Furthermore, the DHP-resistant secretion evoked by KCl was completely blocked by 10  $\mu$ M Cd<sup>2+</sup> (Fig. 5A), a nonselective Ca<sub>v</sub> blocker (Hille, 1995), and by 1  $\mu$ M  $\omega$ -agatoxin IVA (Mintz et al., 1992), a specific Ca<sub>v</sub>2.1 blocker (Fig. 5B).

## Discussion

Ca. 1.3 Is Preferentially Linked to Glucose-Stimulated Insulin Secretion in INS-1 Cells. By using INS-1 cells that stably express DHP-insensitive Ca<sub>v</sub>1.2 or Ca<sub>v</sub>1.3 channels, we were able to pharmacologically isolate these two distinct channels subtypes and to study the contribution of each to insulin secretion. The Ca. 1.2 (rat brain; Snutch et al., 1991) and Ca, 1.3 (human brain; Williams et al., 1992) clones used in this study are nearly identical to the respective channel subtypes isolated from human pancreatic islets (Seino et al., 1992). Electrophysiological characterization of both Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cells demonstrated that the mutant channels were functionally expressed, and that total Ca<sup>2+</sup> channel activity was not grossly different from untransfected INS-1 cells (Fig. 2). Because only the poreforming  $\alpha_1$  subunits were used in the construction of the Ca. 1.2/DHPi and Ca. 1.3/DHPi cell lines, it is likely that the endogenous auxiliary channel subunits were limiting, and thus controlled the number of  $Ca_v \alpha_1$  subunits expressed at the cell surface. Using this model system, we found that both channel subtypes were able to mediate DHP-resistant insu-



Fig. 3. Insulin secretion in Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cells. A, glucose- (11 mM) and KCl (50 mM)-stimulated secretion in INS-1 cells. In untransfected INS-1 cells, 0.1  $\mu$ M PN200-110 completely blocked glucose- or KCl-induced insulin secretion [glucose: basal = 0.38 ± 0.14% (*n* = 3); 11 mM glucose = 1.38 ± 0.12% (*n* = 3) KCl: basal = 4.06 ± 0.63% (*n* = 3); 50 mM KCl = 11.18 ± 1.15% (*n* = 3)] (\*, *p* < 0.05 compared with basal). B, glucose-stimulated secretion in Ca<sub>v</sub>1.2/DHPi or Ca<sub>v</sub>1.3/DHPi cells (11 mM; Glu). Ca<sub>v</sub>1.2/DHPi is basal = 0.99 ± 0.16% (*n* = 3); Glu = 4.06 ± 0.55% (*n* = 3). PN200-110 (1  $\mu$ M) completely inhibited glucose stimulated insulin secretion in Ca<sub>v</sub>1.2/DHPi cells (31 mM; Glu). Ca<sub>v</sub>1.2/DHPi cells (Glu + PN). With Ca<sub>v</sub>1.3/DHPi cells, a significant fraction of glucose-stimulated insulin secretion was resistant to 1  $\mu$ M PN200-110, but blocked by 500  $\mu$ M diltiazem (Glu + PN + Dil). Ca<sub>v</sub>1.3/DHPi cells and Ca<sub>v</sub>1.3/DHPi cells and Ca<sub>v</sub>1.3/DHPi cells (*n* = 3); Glu + PN + Dil = 1.63 ± 0.3% (*n* = 3) (\*, *p* < 0.05 compared with basal). C, KCl-stimulated secretion. Both Ca<sub>v</sub>1.2/DHPi cells and Ca<sub>v</sub>1.3/DHPi cells demonstrated PN200-110-resistant insulin secretion in response to KCl stimulation. Ca<sub>v</sub>1.2/DHPi: basal = 0.83 ± 0.6% (*n* = 3); KCl = 24.3 ± 5% (*n* = 3); KCl + PN = 6.28 ± 0.6% (*n* = 3); KCl + PN + Dil = 1.91 ± 0.9% (*n* = 3) (\*, *p* < 0.05 compared with basal). D, DHP-insensitive insulin secretion in Ca<sub>v</sub>1.3/DHPi cells. In Ca<sub>v</sub>1.3/DHPi cells (*n* = 3); KCl = 0.05% (*n* = 3); KCl + PN + Dil = 2.3 ± 0.1% (*n* = 3) (\*, *p* < 0.05 compared with basal). D, DHP-insensitive insulin secretion in Ca<sub>v</sub>1.3/DHPi cells. In Ca<sub>v</sub>1.3/DHPi cells (*n* = 3); than Ca<sub>v</sub>1.3/DHPi cells. In Ca<sub>v</sub>1.3/DHPi cells (*n* = 3) of the glucose-insulin secretion was resistant to 1  $\mu$ M PN200-110, whereas no DHP-insensitive insulin secretion was detected in Ca<sub>v</sub>1.3/DHPi cells (*n* = 3, \*\*, *p* < 0.01). With KCl stimulation, Ca<sub>v</sub>1.3/DHPi supported a significantly higher level of DHP-insensitive

lin secretion in response to KCl depolarization, whereas only Ca. 1.3 was able to mediate DHP-resistant secretion in response to glucose. Expression of the drug-resistant mutants did not seem to grossly disrupt the endogenous secretory pathway. In both Ca, 1.2/DHPi and Ca, 1.3/DHPi cells, the majority of secretion was blocked by low concentrations of PN200-110, indicating the contribution of endogenous L-type channels to excitation-secretion coupling. A previous study attempted to assess the roles of both Ca<sub>v</sub>1.2 and 1.3 channels in mediating insulin secretion using Cav1.3 knockout mice (Platzer et al., 2000), but that study did not indicate a role for Ca<sub>v</sub>1.3 in insulin secretion. However, a subsequent article suggested that the pancreatic  $\beta$ -cells of the mouse strain used to generate the Ca, 1.3 knockouts did not express Ca, 1.3 (Barg et al., 2001). Understanding how Ca<sub>v</sub>1.3 is coupled to insulin secretion is of therapeutic interest because the predominant  $Ca^{2+}$  channel in human pancreatic  $\beta$  cells is  $Ca_{..}1.3$ (Seino et al., 1992). This study also suggests that drug-insensitive channels may facilitate study of channel regulation or other channel-mediated events in the native cell type or tissue.

Intracellular II-III Loop of Ca<sub>v</sub>1.3 Plays a Critical Role in Glucose-Stimulated Insulin Secretion. We further demonstrated the critical role of Ca, 1.3 in glucose-stimulated insulin secretion using INS-1 cells that stably express the intracellular II-III loop of either Ca, 1.2 or Ca, 1.3. Ca, 1.2/ II-III cells were not different from untransfected INS-1 cells in both KCl- and glucose-stimulated insulin secretion. However, the endogenous L-type channels in Ca. 1.3/II-III cells seemed to be largely uncoupled from insulin secretion. This uncoupling completely abolished insulin secretion in response to glucose, and markedly reduced KCl-stimulated secretion. The observation that 1  $\mu$ M  $\omega$ -agatoxin IVA could inhibit the DHP-resistant fraction of KCl-stimulated insulin secretion in these cells suggests that the Ca<sub>2</sub>.1 channels endogenous to INS-1 cells can effectively couple to KCl- but not glucose-stimulated insulin secretion. However, the endogenous L-type channels in Ca, 1.3/II-III cell are functional because in the presence of FPL 64176, KCl-stimulated secretion is restored to levels similar to untransfected INS-1 cells. We speculate that the increased  $Ca^{2+}$  influx induced by FPL 64176 compensates for a greater distance between the chan-



**Fig. 4.** Insulin secretion in Ca<sub>v</sub>1.2/II-III and Ca<sub>v</sub>1.3/II-III cells. A, intracellular II-III loop of Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3. Both Ca<sub>v</sub>1.2/II-III and Ca<sub>v</sub>1.3/II-III were subcloned into the EGFP vector such that GFP was fused to the C terminus of each and stably transfected into INS-1 cells. B, stable expression of Ca<sub>v</sub>1.2 II-III or Ca<sub>v</sub>1.3 II-III in INS-1 cells. Western blot of whole-cell lysates with anti-GFP antibodies detected proteins of 43 and 45 kDa corresponding to Ca<sub>v</sub>1.3 II-III/GFP and Ca<sub>v</sub>1.2 II-III/GFP, respectively. The lower molecular mass band from Ca<sub>v</sub>1.3/II-III cells most likely represents a small amount of proteolytic cleavage of the fusion to generate a fragment containing GFP and little, if any, of the Ca<sub>v</sub>1.3 II-III loop. C, glucose (11 mM; Glu) did not stimulate insulin secretion in Ca<sub>v</sub>1.3/II-III cells (n = 3). In Cav1.2/II-III cells, glucose-stimulated insulin secretion was similar to untransfected cells (basal =  $1.23 \pm 0.15\%$ ; Glu =  $2.2 \pm 0.08\%$ ) and completely blocked by 0.1  $\mu$ M PN200-110 (PN) (n = 3). D, both Ca<sub>v</sub>1.2/II-III and Ca<sub>v</sub>1.3/II-III cells was blocked by 0.1  $\mu$ M PN200-110 (PN) (basal =  $5.33 \pm 0.24\%$ ; KCl =  $11.4 \pm 0.7\%$ ; KCl + PN =  $6.51 \pm 0.63\%$  (n = 3)). KCl-stimulated secretion in Ca<sub>v</sub>1.3/II-III cells was resistant to 0.1  $\mu$ M (0.1 PN) and 1.0  $\mu$ M PN200-110 (1PN) [basal =  $1.03 \pm 0.09\%$ ; KCl =  $4.42 \pm 0.32\%$ ; KCl +  $0.1PN = 3.7 \pm 0.21\%$ ; KCl + 1PN =  $3.16 \pm 0.57\%$  (n = 3)] (\*, p < 0.05 compared with basal).

nel and the Ca<sup>2+</sup> sensor of the secretory machinery. A similar mechanism was proposed to explain inhibition of evoked neurotransmitter release upon injection of a peptide corresponding to the II-III loop of Ca<sub>v</sub>2.2 into the presynaptic cell of a cultured synapse preparation (Rettig et al., 1997). This inhibition was reversed by increasing the extracellular Ca<sup>2+</sup> concentration. This result suggests that the intracellular II-III loop of Ca<sub>v</sub>1.3 plays a prominent role in linking the Ca<sub>v</sub>1.3 channel specifically to glucose-stimulated secretion, and that overexpressed Ca, 1.3 II-III loop competitively inhibits a key protein-protein interaction. A binding partner for the Cav1.3 II-III loop has not yet been clearly identified. Syntaxin has been shown to interact with the II-III loop of Ca, 1.2 (Wiser et al., 1999) and to colocalize with Ca. 1.3 channels in pancreatic  $\beta$ -cells (Yang et al., 1999). However, neither of these studies specifically examined the role of syntaxin/L-type channel interactions in glucose-stimulated insulin secretion. Because we found that Ca. 1.3, but not Ca. 1.2, can contribute to glucose-induced secretion in INS-1 cells, our results suggest that L-type channel/syntaxin interactions may not be sufficient to efficiently couple glucose-induced membrane depolarization to insulin secretion. The II-III loops of the channels used in this study are only 43% identical, and the II-III loop of Ca, 1.3 contains two consensus protein kinase A phosphorylation sites not present in Ca, 1.2. In addition, the Ca, 1.3 II-III loop contains two potential SH-3 domain binding sites (PXXP motifs; Mayer, 2001) that are not conserved in Ca, 1.2. Identification of the protein(s) that interacts with the II-III loop of  $\rm Ca_v 1.3$  to mediate this specific coupling will give further insight into the mechanism of glucose-stimulated insulin secretion.

Potential Mechanism for Specificity of Ca. 1.3 Coupling to Insulin Secretion. Although interactions between SNARE proteins and voltage-gated Ca<sup>2+</sup> channels are clearly important for vesicular exocytosis of neurotransmitters, it seems unlikely that such interactions alone could account for the preferential linkage of Ca. 1.3 to glucose-stimulated insulin secretion that we observed. Because glucose induces smaller depolarization of membrane potential in INS-1 cells than 30 mM KCl (Kennedy et al., 1998; Antunes et al., 2000), one potential explanation for our results is that Ca, 1.3 channels are activated at more negative potentials than Ca. 1.2 channels. Several studies have concluded that various splice variants of Ca<sub>v</sub>1.3 activate at more negative potentials than Cav1.2 (Koschak et al., 2001; Scholze et al., 2001; Xu and Lipscombe, 2001). However, the voltage dependence of activation for the Ca<sub>v</sub>1.3 and Ca<sub>v</sub>1.2 clones used in this study are virtually identical when measured in human embryonic kidney 293 cells (Bell et al., 2001; Gage et al., 2002). Furthermore, we have not observed any significant difference in the current-voltage relationship of the DHPi fraction of current between the Ca<sub>v</sub>1.2/DHPi and Ca<sub>v</sub>1.3/DHPi cell lines using conventional whole recordings (G. Liu and G. H. Hockerman, unpublished observations). However, we cannot rule out the possibility that glucose metabolism by intact INS-1 cells can shift the activation threshold of Cav1.3 to more negative



Fig. 5. Endogenous L-type Ca<sup>2+</sup> channels are functional in Ca<sub>v</sub>1.3/II-III cells. A, stimulation with 50 mM KCl (KCl) induces insulin secretion from Ca<sub>v</sub>1.3/II-III cells, which is largely resistant to 1  $\mu$ M PN200-110 (PN). FPL 64176 (10  $\mu$ M; KCl + FPL) markedly potentiated KCl-induced insulin secretion in Ca<sub>v</sub>1.3/II-III cells. The potentiation was blocked by 1  $\mu$ M PN200-110 (KCl + FPL + PN). The DHP-resistant insulin secretion was completely blocked by 10  $\mu$ M Cd<sup>2+</sup> (KCl + Cd<sup>2+</sup>) (\*, p < 0.05 compared with basal). Inset, The increase in KCl-stimulated insulin secretion with 0  $\mu$ M FPL 64176 in Ca<sub>v</sub>1.3/II-III cells was greater (443 ± 1.5%) than in untransfected INS-1 cells (174 ± 25%) (\*, p < 0.05 compared with INS-1 cells). B, PN200-110 (1  $\mu$ M) blocked virtually all KCl-stimulated insulin secretion in Ca<sub>v</sub>1.3/II-III cells (PN; \*, p < 0.05 compared insulin secretion in INS-1 cells (INS-1 + PN; 95 ± 9.3%), but only 42.5 ± 5.5% of KCl-stimulated secretion in Ca<sub>v</sub>1.3/II-III cells (PN; \*, p < 0.05). However, addition of both 1  $\mu$ M PN200-110 and 1  $\mu$ M  $\omega$ -agatoxin IVA to Ca<sub>v</sub>1.3/II-III cells completely blocked KCl-stimulated secretion (PN +  $\omega$ -Aga; 96.4 ± 6.3).

potentials (Smith et al., 1989). Therefore, it is unlikely that differences in  $Ca_v 1.2$  and  $Ca_v 1.3$  channel activation threshold can explain our results unless this property of these channels is differentially modulated in intact INS-1 cells.

An alternative explanation for our results is that Ca<sup>2+</sup> influx via Ca, 1.3 channels is specifically coupled to an intracellular signaling mechanism that is not activated by Ca<sup>2+</sup> influx via Ca<sub>v</sub>1.2. Although INS-1 cells retain glucose-induced insulin secretion, RINm5F cells, another rat  $\beta$ -cell line, do not, although both cell lines express Ca. 1.3 (Safayhi et al., 1997; Horvath et al., 1998). Interestingly, RINm5F cells do not express the endoplasmic reticulum Ca<sup>2+</sup> release channel RYR2, whereas INS-1 cells do (Gamberucci et al., 1999). The role of Ca<sup>2+</sup> release from internal stores in insulin secretion is not clear. However, a recent report concluded that L-type Ca<sup>2+</sup> channel activity initiates Ca<sup>2+</sup> induced  $Ca^{2+}$  release via RYR2 in mouse  $\beta$ -cells (Lemmens et al., 2001). The L-type Ca<sup>2+</sup> channel from skeletal muscle, Ca, 1.1, is coupled to RYR1 in the sarcoplasmic reticulum, via the intracellular II-III loop (Grabner et al., 1999). Thus, it is attractive to speculate that a similar coupling between  $Ca_v 1.3$  and intracellular  $Ca^{2+}$  release may mediate a specific role for  $Ca_v 1.3$  in glucose-stimulated insulin secretion.

In summary, we have shown that  $Ca_v 1.3$  is preferentially linked to glucose-stimulated insulin secretion in INS-1 cells and that the intracellular loop between homologous domains II and III is likely involved in mediating this specificity. It will be of interest to investigate the mechanism for this specificity, and specifically, to determine whether the II-III loop of  $Ca_v 1.3$  interacts with other proteins in this process.

#### Acknowledgments

PN200-110 was the gift of Novartis (Basel, Switzerland). The human brain Ca<sub>v</sub>1.3 clone was provided by SIBIA Neurosciences (San Diego, CA), now Merck Research Laboratories. We thank Dr. Ming Li for the gift of INS-1 cells and helpful discussions regarding their culture.

#### References

- Antunes CM, Salgado AP, Rosario LM, and Santos RM (2000) Differential patterns of glucose-induced electrical activity and intracellular calcium responses in single mouse and rat pancreatic islets. *Diabetes* 49:2028–2038.
- Asfari M, Janjic D, Meda P, Li G, Halban PA, and Wollheim CB (1992) Establishment of 2-mercaptoethanol-dependent differentiated insulin-secreting cell lines. *Endocrinology* 130:167-178.
- Barg S, Ma XS, Eliasson L, Galvanovskis J, Gopel SO, Obermuller S, Platzer J, Renstrom E, Trus M, Atlas D, et al. (2001) Fast exocytosis with few Ca<sup>2+</sup> channels in insulin-secreting mouse pancreatic B cells. *Biophys J* 81:3308–3323.
- Bell DC, Butcher AJ, Berrow NS, Page KM, Brust PF, Nesterova A, Stauderman KA, Seabrook GR, Nurnberg B, and Dolphin AC (2001) Biophysical properties, pharmacology and modulation of human, neuronal L-type (α(1D), Ca(V)1.3) voltagedependent calcium currents. J Neurophysiol 85:816-827.
- Bokvist K, Eliasson L, Ammala C, Renstrom E, and Rorsman P (1995) Colocalization of L-type  $Ca^{2+}$  channels and insulin-containing secretory granules and its significance for the initiation of exocytosis in mouse pancreatic B-cells. *EMBO* (*Eu Mol Bio Organ*) J 14:50-57.
- Devis G, Somers G, Van Obberghen E, and Malaisse WJ (1975) Calcium antagonists and islet function. I. Inhibition of insulin release by verapamil. *Diabetes* 24:247– 251.
- Gage MJ, Rane SG, Hockerman GH, and Smith TJ (2002) The virally encoded fungal toxin KP4 specifically blocks L-type voltage-gated calcium channels. *Mol Pharma-* col **61**:936–944.
- Gamberucci A, Fulceri R, Pralong W, Banhegyi G, Marcolongo P, Watkins SL, and Benedetti A (1999) Caffeine releases a glucose-primed endoplasmic reticulum Ca<sup>2+</sup> pool in the insulin secreting cell line INS-1. *FEBS Lett* **446**:309–312.
- Grabner M, Dirksen RT, Suda N, and Beam KG (1999) The ii-iii loop of the skeletal muscle dihydropyridine receptor is responsible for the bi-directional coupling with the ryanodine receptor. J Biol Chem 274:21913–21919.
- Hille B (1995) Ionic Channels of Excitable Membranes. Sinauer, Sunderland, MA. Hockerman GH, Dilmac N, Scheuer T, and Catterall WA (2000) Molecular determinants of diltiazem block in domains IIIS6 and IVS6 of L-type Ca<sup>2+</sup> channels. Mol Pharmacol 58:1264–1270.
- Hockerman GH, Peterson BZ, Johnson BD, and Catterall WA (1997) Molecular determinants of drug binding and action on L-type Ca<sup>2+</sup> channels. Annu Rev Pharmacol Toxicol 37:361–396.
- Horvath A, Szabadkai G, Varnai P, Aranyi T, Wollheim CB, Spat A, and Enyedi P (1998) Voltage dependent calcium channels in adrenal glomerulosa cells and in insulin producing cells. *Cell Calcium* 23:33–42.
- Jacobsson G, Bean AJ, Scheller RH, Juntti-Berggren L, Deeney JT, Berggren PO, and Meister B (1994) Identification of synaptic proteins and their isoform mRNAs in compartments of pancreatic endocrine cells. *Proc Natl Acad Sci USA* 91:12487– 12491.
- Kennedy ED, Maechler P, and Wollheim CB (1998) Effects of depletion of mitochondrial DNA in metabolism secretion coupling in INS-1 cells. *Diabetes* 47:374–380.
- Koschak A, Reimer D, Huber I, Grabner M, Glossmann H, Engel A, and Striessnig J (2001)  $\alpha_{1D}$  (Ca<sub>v</sub>1.3) subunits can form L-type channels activating at negative voltages. J Biol Chem **276:**22100–22106.
- Lemmens R, Larsson O, Berggren PO, and Islam MS (2001)  $Ca^{2+}$ -induced  $Ca^{2+}$  release from the endoplasmic reticulum amplifies the  $Ca^{2+}$  signal mediated by activation of voltage-gated L-type  $Ca^{2+}$  channels in pancreatic  $\beta$ -cells. *J Biol Chem* **276**:9971–9977.
- Ligon B, Boyd AE 3rd, and Dunlap K (1998) Class a calcium channel variants in pancreatic islets and their role in insulin secretion. *J Biol Chem* **273**:13905–13911.
- Mayer BJ (2001) SH3 domains: complexity in moderation. *J Cell Sci* **114**:1253–1263. Mintz IM, Venema VJ, Swiderek KM, Lee TD, Bean BP, and Adams ME (1992)
- P-type calcium channels blocked by the spider toxin omega-Aga-IVA. *Nature* (Lond) **355**:827–829.
- Nagamatsu S, Fujiwara T, Nakamichi Y, Watanabe T, Katahira H, Sawa H, and Akagawa K (1996) Expression and functional role of syntaxin 1/hpc-1 in pancreatic  $\beta$  cells. Syntaxin 1a, but not 1b, plays a negative role in regulatory insulin release pathway. J Biol Chem **271**:1160–1165.
- Peterson BZ, Johnson BD, Hockerman GH, Acheson M, Scheuer T, and Catterall WA

(1997) Analysis of the dihydropyridine receptor site of l-type calcium channels by

- alanine-scanning mutagenesis. J Biol Chem **272:**18752–18758. Platzer J, Engel J, Schrott-Fischer A, Stephan K, Bova S, Chen H, Zheng H, and Striessnig J (2000) Congenital deafness and sinoatrial node dysfunction in mice lacking class d L-type Ca2+ channels. Cell 102:89-97.
- Qian WJ and Kennedy RT (2001) Spatial organization of Ca(2+) entry and exocytosis in mouse pancreatic  $\beta$ -cells. Biochem Biophys Res Commun 286:315-321.
- Rajan AS, Aguilar-Bryan L, Nelson DA, Yaney GC, Hsu WH, Kunze DL, and Boyd AE 3d. (1990) Ion channels and insulin secretion. Diabetes Care 13:340-363.
- Rettig J, Heinemann C, Ashery U, Sheng ZH, Yokoyama CT, Catterall WA, and Neher E (1997) Alteration of  $Ca^{2+}$  dependence of neurotransmitter release by disruption of  $Ca^{2+}$  channel/syntaxin interaction. J Neurosci 17:6647–6656.
- Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, and Catterall WA (1996) Isoform-specific interaction of the  $\alpha$ 1a subunits of brain Ca<sup>2+</sup> channels with the presynaptic proteins syntaxin and SNAP-25. Proc Natl Acad Sci USA 93:7363-7368.
- Sadoul K, Lang J, Montecucco C, Weller U, Regazzi R, Catsicas S, Wollheim CB, and Halban PA (1995) SNAP-25 is expressed in islets of Langerhans and is involved in insulin release. J Cell Biol 128:1019-1028.
- Safavhi H, Haase H, Kramer U, Bihlmaver A, Roenfeldt M, Ammon HP, Froschmavr M, Cassidy TN, Morano I, Ahlijanian MK, et al. (1997) L-Type calcium channels in insulin-secreting cells: biochemical characterization and phosphorylation in rinm5f cells. Mol Endocrinol 11:619-629.
- Scholze A, Plant TD, Dolphin AC, and Nurnberg B (2001) Functional expression and characterization of a voltage-gated  $Ca_v 1.3 (\alpha_{1D})$  calcium channel subunit from an insulin-secreting cell line. Mol Endocrinol 15:1211-1221.
- Seino S, Chen L, Seino M, Blondel O, Takeda J, Johnson JH, and Bell GI (1992) Cloning of the  $\alpha 1$  subunit of a voltage-dependent calcium channel expressed in pancreatic β cells. Proc Natl Acad Sci USÂ 89:584-588.
- Sheng ZH, Rettig J, Takahashi M, and Catterall WA (1994) Identification of a syntaxin-binding site on N-type calcium channels. Neuron 13:1303-1313.

- Sheng ZH, Westenbroek RE, and Catterall WA (1998) Physical link and functional coupling of presynaptic calcium channels and the synaptic vesicle docking/fusion machinery. J Bioenerg Biomem 30:335-345.
- Smith PA, Rorsman P, and Ashcroft FM (1989) Modulation of dihydropyridinesensitive  $Ca^{2+}$  channels by glucose metabolism in mouse pancreatic  $\beta$ -cells. Nature (Lond) 342:550-553.
- Snutch TP, Tomlinson WJ, Leonard JP, and Gilbert MM (1991) Distinct calcium channels are generated by alternative splicing and are differentially expressed in the mammalian cns. Neuron 7:45-57.
- Williams ME, Feldman DH, McCue AF, Brenner R, Velicelebi G, Ellis SB, and Harpold MM (1992) Structure and functional expression of  $\alpha 1$ ,  $\alpha 2$  and  $\beta$  subunits of a novel human neuronal calcium channel subtype. Neuron 8:71-84.
- Wiser O, Trus M, Hernandez A, Renstrom E, Barg S, Rorsman P, and Atlas D (1999) The voltage sensitive LC-type Ca<sup>2+</sup> channel is functionally coupled to the exocytotic machinery. Proc Natl Acad Sci USA 96:248-253.
- Wollheim CB and Sharp GW (1981) Regulation of insulin release by calcium. Physiol Rev 61:914-973.
- Xu W and Lipscombe D (2001) Neuronal  $Ca_v 1.3\alpha_1$  L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. J Neurosci 21:5944-5951.
- Yang S-N, Larsson O, Branstrom R, Bertorello A, Leibiger B, Leibiger IB, Moede T, Kohler M, Meister B, and Berggren P-O (1999) Syntaxin 1 interacts with the LD subtype of voltage-gated  $Ca^{2+}$  channels in pancreatic  $\beta$  cells. Proc Natl Acad Sci USA 96:10164-10169.
- Zhuang H, Bhattacharjee A, Hu F, Zhang M, Goswami T, Wang L, Wu S, Berggren PO, and Li M (2000) Cloning of a T-type Ca<sup>2+</sup> channel isoform in insulin-secreting cells. Diabetes 49:59-64.

Address correspondence to: Gregory H. Hockerman, 575 Stadium Mall Dr., West Lafayette, IN 47907-209. E-mail: gregh@pharmacy.purdue.edu